


# PALS Pulseless Arrest Algorithm.



**Doses/Details**

**CPR Quality**

- Push hard ( $\geq 1/3$  of anterior-posterior diameter of chest) and fast (at least 100/min) and allow complete chest recoil
- Minimize interruptions in compressions
- Avoid excessive ventilation
- Rotate compressor every 2 minutes
- If no advanced airway, 15:2 compression-ventilation ratio. If advanced airway, 8-10 breaths per minute with continuous chest compressions

**Shock Energy for Defibrillation**

First shock 2 J/kg, second shock 4 J/kg, subsequent shocks  $\geq 4$  J/kg, maximum 10 J/kg or adult dose.

**Drug Therapy**

- **Epinephrine IO/IV Dose:** 0.01 mg/kg (0.1 mL/kg of 1:10 000 concentration). Repeat every 3-5 minutes. If no IO/IV access, may give endotracheal dose: 0.1 mg/kg (0.1 mL/kg of 1:1000 concentration).
- **Amiodarone IO/IV Dose:** 5 mg/kg bolus during cardiac arrest. May repeat up to 2 times for refractory VF/pulseless VT.

**Advanced Airway**

- Endotracheal intubation or supraglottic advanced airway
- Waveform capnography or capnometry to confirm and monitor ET tube placement
- Once advanced airway in place give 1 breath every 6-8 seconds (8-10 breaths per minute)

**Return of Spontaneous Circulation (ROSC)**

- Pulse and blood pressure
- Spontaneous arterial pressure waves with intra-arterial monitoring


**Reversible Causes**

- Hypovolemia
- Hypoxia
- Hydrogen ion (acidosis)
- Hypoglycemia
- Hypo-/hyperkalemia
- Hypothermia
- Tension pneumothorax
- Tamponade, cardiac
- Toxins
- Thrombosis, pulmonary
- Thrombosis, coronary

Kleinman M E et al. Circulation 2010;122:S876-S908

# PALS Bradycardia Algorithm.

## Pediatric Bradycardia With a Pulse and Poor Perfusion




© 2010 American Heart Association

Kleinman M E et al. Circulation 2010;122:S876-S908

# PALS Tachycardia Algorithm.

## Pediatric Tachycardia With a Pulse and Poor Perfusion



Kleinman M E et al. Circulation 2010;122:S876-S908

# Recognition of Respiratory Problems Flowchart



<b>Pediatric Advanced Life Support</b> <b>Signs of Respiratory Problems</b>					
Clinical Signs		Upper Airway Obstruction	Lower Airway Obstruction	Lung Tissue Disease	Disordered Control of Breathing
<b>A</b>	Patency	Airway open and maintainable/not maintainable			
<b>B</b>	Respiratory Rate/Effort	Increased			Variable
	Breath Sounds	Stridor (typically inspiratory) Barking cough Hoarseness	Wheezing (typically expiratory) Prolonged expiratory phase	Grunting Crackles Decreased breath sounds	Normal
	Air Movement	Decreased			Variable
<b>C</b>	Heart Rate	Tachycardia (early)		Bradycardia (late)	
	Skin	Pallor, cool skin (early)		Cyanosis (late)	
<b>D</b>	Level of Consciousness	Anxiety, agitation (early) Lethargy, unresponsiveness (late)			
<b>E</b>	Temperature	Variable			
<b>Pediatric Advanced Life Support</b> <b>Identification of Respiratory Problems by Severity</b>					
<div style="display: flex; justify-content: space-between; align-items: center;"> <div style="text-align: center;"> <b>Respiratory Distress</b> </div> <div style="text-align: center;"> <b>Respiratory Failure</b> </div> </div>					
<b>A</b>	Open and maintainable → <b>Not maintainable</b>				
<b>B</b>	Tachypnea → <b>Bradypnea to apnea</b>				
	Work of breathing (nasal flaring/retractions) <b>Increased effort</b> → <b>Decreased effort</b> → <b>Apnea</b>				
	Good air movement → <b>Poor to absent air movement</b>				
<b>C</b>	Tachycardia → <b>Bradycardia</b>				
	Pallor → <b>Cyanosis</b>				
<b>D</b>	Anxiety, agitation → <b>Lethargy to unresponsiveness</b>				
<b>E</b>	Variable temperature				

# Management of Shock Flowchart



<b>Management of Shock Flowchart</b>			
<ul style="list-style-type: none"> <li>• Oxygen</li> <li>• Pulse oximetry</li> <li>• ECG monitor</li> </ul>		<ul style="list-style-type: none"> <li>• IV/IO access</li> <li>• BLS as indicated</li> <li>• Point-of-care glucose testing</li> </ul>	
<b>Hypovolemic Shock</b>			
<b>Specific Management for Selected Conditions</b>			
<b>Nonhemorrhagic</b>		<b>Hemorrhagic</b>	
<ul style="list-style-type: none"> <li>• 20 mL/kg NS/LR bolus, repeat as needed</li> <li>• Consider colloid</li> </ul>		<ul style="list-style-type: none"> <li>• Control external bleeding</li> <li>• 20 mL/kg NS/LR bolus, repeat 2 or 3× as needed</li> <li>• Transfuse PRBCs as indicated</li> </ul>	
<b>Distributive Shock</b>			
<b>Specific Management for Selected Conditions</b>			
<b>Septic</b>	<b>Anaphylactic</b>		<b>Neurogenic</b>
Management Algorithm: <ul style="list-style-type: none"> <li>• Septic Shock</li> </ul>	<ul style="list-style-type: none"> <li>• IM epinephrine (or autoinjector)</li> <li>• Fluid boluses (20 mL/kg NS/LR)</li> <li>• Albuterol</li> <li>• Antihistamines, corticosteroids</li> <li>• Epinephrine infusion</li> </ul>		<ul style="list-style-type: none"> <li>• 20 mL/kg NS/LR bolus, repeat PRN</li> <li>• Vasopressor</li> </ul>
<b>Cardiogenic Shock</b>			
<b>Specific Management for Selected Conditions</b>			
<b>Bradycardia/Tachycardia</b>		<b>Other (eg, CHD, Myocarditis, Cardiomyopathy, Poisoning)</b>	
Management Algorithms: <ul style="list-style-type: none"> <li>• Bradycardia</li> <li>• Tachycardia With Poor Perfusion</li> </ul>		<ul style="list-style-type: none"> <li>• 5 to 10 mL/kg NS/LR bolus, repeat PRN</li> <li>• Vasoactive infusion</li> <li>• Consider expert consultation</li> </ul>	
<b>Obstructive Shock</b>			
<b>Specific Management for Selected Conditions</b>			
<b>Ductal-Dependent (LV Outflow Obstruction)</b>	<b>Tension Pneumothorax</b>	<b>Cardiac Tamponade</b>	<b>Pulmonary Embolism</b>
<ul style="list-style-type: none"> <li>• Prostaglandin E<sub>1</sub></li> <li>• Expert consultation</li> </ul>	<ul style="list-style-type: none"> <li>• Needle decompression</li> <li>• Tube thoracostomy</li> </ul>	<ul style="list-style-type: none"> <li>• Pericardiocentesis</li> <li>• 20 mL/kg NS/LR bolus</li> </ul>	<ul style="list-style-type: none"> <li>• 20 mL/kg NS/LR bolus, repeat PRN</li> <li>• Consider thrombolytics, anticoagulants</li> <li>• Expert consultation</li> </ul>

## Recognition of Shock Flowchart



Clinical Signs		Hypovolemic Shock	Distributive Shock	Cardiogenic Shock	Obstructive Shock
<b>A</b>	Patency	Airway open and maintainable/not maintainable			
<b>B</b>	Respiratory rate	Increased			
	Respiratory effort	Normal to increased		Labored	
	Breath sounds	Normal	Normal (± crackles)	Crackles, grunting	
<b>C</b>	Systolic blood pressure	<b>Compensated Shock → Hypotensive Shock</b>			
	Pulse pressure	Narrow	Variable	Narrow	
	Heart rate	Increased			
	Peripheral pulse quality	Weak	Bounding or weak	Weak	
	Skin	Pale, cool	Warm or cool	Pale, cool	
	Capillary refill	Delayed	Variable	Delayed	
	Urine output	Decreased			
<b>D</b>	Level of consciousness	Irritable early Lethargic late			
<b>E</b>	Temperature	Variable			

# Management of Respiratory Emergencies Flowchart



## Management of Respiratory Emergencies Flowchart

- Airway positioning
- Suction as needed
- Oxygen
- Pulse oximetry
- ECG monitor (as indicated)
- BLS as indicated

### Upper Airway Obstruction Specific Management for Selected Conditions

Croup	Anaphylaxis	Aspiration Foreign Body
<ul style="list-style-type: none"> <li>• Nebulized epinephrine</li> <li>• Corticosteroids</li> </ul>	<ul style="list-style-type: none"> <li>• IM epinephrine (or autoinjector)</li> <li>• Albuterol</li> <li>• Antihistamines</li> <li>• Corticosteroids</li> </ul>	<ul style="list-style-type: none"> <li>• Allow position of comfort</li> <li>• Specialty consultation</li> </ul>

### Lower Airway Obstruction Specific Management for Selected Conditions

Bronchiolitis	Asthma
<ul style="list-style-type: none"> <li>• Nasal suctioning</li> <li>• Bronchodilator trial</li> </ul>	<ul style="list-style-type: none"> <li>• Albuterol ± ipratropium</li> <li>• Corticosteroids</li> <li>• Subcutaneous epinephrine</li> <li>• Magnesium sulfate</li> <li>• Terbutaline</li> </ul>

### Lung Tissue Disease Specific Management for Selected Conditions

Pneumonia/Pneumonitis			Pulmonary Edema	
Infectious	Chemical	Aspiration	Cardiogenic or Noncardiogenic (ARDS)	
<ul style="list-style-type: none"> <li>• Albuterol</li> <li>• Antibiotics (as indicated)</li> </ul>			<ul style="list-style-type: none"> <li>• Consider noninvasive or invasive ventilatory support with PEEP</li> <li>• Consider vasoactive support</li> <li>• Consider diuretic</li> </ul>	

### Disordered Control of Breathing Specific Management for Selected Conditions

Increased ICP	Poisoning/Overdose	Neuromuscular Disease
<ul style="list-style-type: none"> <li>• Avoid hypoxemia</li> <li>• Avoid hypercarbia</li> <li>• Avoid hyperthermia</li> </ul>	<ul style="list-style-type: none"> <li>• Antidote (if available)</li> <li>• Contact poison control</li> </ul>	<ul style="list-style-type: none"> <li>• Consider noninvasive or invasive ventilatory support</li> </ul>